

A Meta-Analysis of Human Acellular Dermis and Submuscular Tissue Expander Breast Reconstruction

Foot

AQ: 1

John Y. S. Kim, M.D.
 Armando A. Davila, B.S.
 Scott Persing, B.A., M.P.H.
 Caitlin M. Connor, B.A.
 Borko Jovanovic, Ph.D.
 Seema Khan, M.D.
 Neil Fine, M.D.
 Vinay Rawlani, M.D.
Chicago and Maywood, Ill.

Background: Human acellular dermal matrix has become an increasingly used adjunct to traditional submuscular tissue expander/implant breast reconstruction, but there is no strong consensus regarding complication outcomes. This study stratified outcomes based on a meta-analysis of complications.

Methods: A query of the MEDLINE database for articles on human acellular dermal matrix and submuscular tissue expander breast reconstruction yielded 901 citations. Two levels of screening identified 48 relevant studies. The DerSimonian and Laird random-effects model was used to perform the meta-analysis. Risk ratios and pooled complication rates were calculated for each outcome of interest.

Results: Nineteen studies reporting human acellular dermal matrix ($n = 2037$) and 35 reporting submuscular outcomes ($n = 12,847$) were used to estimate complication rates. Rates were generally higher in acellular dermis patients: total complications, 15.4 versus 14.0 percent; seroma, 4.8 versus 3.5 percent; infection, 5.3 versus 4.7 percent; and flap necrosis, 6.9 versus 4.9 percent. Six studies reporting both acellular dermis and submuscular outcomes were used to estimate relative risks. There was an increased risk of total complications (relative risk, 2.05; 95 percent CI, 1.55 to 2.70), seroma (relative risk, 2.73; 95 percent CI, 1.67 to 4.46), infection (relative risk, 2.47; 95 percent CI, 1.71 to 3.57), and reconstructive failure (relative risk, 2.80; 95 percent CI, 1.76 to 4.45) in acellular dermis patients.

Conclusions: The meta-analysis suggests that the use of human acellular dermal matrix increases complication rates vis-à-vis submuscular expander/implant reconstruction. This must be weighed against its reported advantages in enhancing cosmesis and ameliorating contracture. (*Plast Reconstr Surg* 129: 1, 2012.)



CLINICAL QUESTION/LEVEL OF EVIDENCE: Therapeutic, III.

In the United States, approximately 57,000 tissue expander/implant-based reconstructions are performed annually, representing approximately 65 percent of all breast reconstructions.¹ Acellular dermis has been an increasingly popular adjunct to traditional expander reconstruction with putative benefits including improved inframammary control, decreased incidence of migration, greater intraoperative fill (with decreased

concomitant expansion time and number of postoperative visits), improved cosmesis, and amelioration of contracture.²⁻⁸ The basic technique involves releasing the pectoralis muscle along its inferior border and using the acellular dermis to reconstruct the ensuing lower pole defect.

The technique of using human acellular dermal matrix for soft-tissue reconstruction has existed since the early 1990s, and its use has been described in a myriad of clinical contexts from

From the Division of Plastic and Reconstructive Surgery, the Department of Preventative Medicine, and the Lynn Sage Breast Center, Northwestern University, Feinberg School of Medicine, and Loyola University Chicago, Stritch School of Medicine.

Received for publication March 21, 2011; accepted June 16, 2011.

Copyright ©2011 by the American Society of Plastic Surgeons
 DOI: 10.1097/PRS.0b013e3182361fde

Disclosure: Dr. Kim is a consultant for and receives research funding from Mentor and the Musculoskeletal Transplant Foundation. Dr. Fine receives research funding from Allergan, Inc. The remaining authors have no financial relationships to disclose.

burn treatment and neurethra reconstruction, to ameliorating contracture in hand and breast surgery.^{2,4,9-11} However, universal acceptance in these varying clinical indications has been tempered by concerns about surgical complications including seromas, infections, and loss of mechanical integrity.¹²⁻¹⁴ The complication profile of human acellular dermal matrix in breast reconstruction ranges widely throughout the 6 years of published reports (range, 3.2 to 48.7 percent), making it difficult to develop generalized outcomes from study to study.^{2,3,5,7,12,15-18} The central unknown remains regarding how outcomes of human acellular dermal matrix–assisted breast reconstruction compare with outcomes of traditional submuscular breast reconstruction. Our study attempts to collate recent studies in *both* treatment arms and determine general complication profiles and patterns by means of standard meta-analysis methodology.

PATIENTS AND METHODS

Search Methods

A literature search was performed using PubMed to query the MEDLINE database (January 1, 2000, to February 6, 2011). Search terms included “tissue expander,” “implant,” “acellular dermis,” “acellular dermal matrix,” “AlloDerm” or “acellular matrix,” and “breast reconstruction.” Additional search methods included a manual review of reference lists of relevant studies.

Selection Criteria

Selection criteria were defined a priori. Eligible studies were limited to English-only and had to examine breast reconstruction in human patients. Study selection underwent two levels of review by two independent researchers (Fig. 1). Each study was required to clearly indicate the number of reconstructions performed. Studies reporting fewer than 25 tissue expander/implant reconstructions in at least one cohort were excluded. Titles and abstracts were screened for the following exclusion criteria: publications of brief communications, correspondence, discussions, letters, conference/lecture manuscripts, case reports, and reviews; publications containing only abstracts; novel modifications of surgical technique; outcomes related to breast augmentation; outcomes related solely to autogenous reconstruction; and outcomes about only a specific high-risk population.

Full articles were then retrieved for all studies that met the first level of criteria. Studies needed

to report or provide data to calculate a total complication rate and report or provide data to calculate at least one of the following postoperative complications: seroma, hematoma, infection, and flap necrosis. In any studies that reported similar or overlapping cohorts, the publication with the greatest number of reconstructions was included.

Data Collection and Analysis

Data collection and analysis was performed following the guidelines set forth by the *Cochrane Handbook for Systematic Reviews of Interventions* and the “Meta-Analysis of Observational Studies in Epidemiology.”^{19,20} Two independent reviewers extracted data from all selected studies by using a standardized data abstraction form. This electronic data form included the lead author, publication year, type of reconstructive procedure, number of patients, number of reconstructions, number of unilateral and bilateral reconstructions, average patient age and body mass index, percentage of smokers and diabetic patients, percentage of patients who received radiation therapy before surgery, percentage of patients who received postoperative radiation therapy, percentage of patients who received chemotherapy, average tissue expander intraoperative fill, and average follow-up. Complication data included number of reconstructions with seromas, hematomas, infection, flap necrosis, and explantation.

All rates used in the analysis were based on the number of reconstructions in each study. Two levels of analysis were performed. Cumulative pooled estimates were calculated from the standard error of complication rates based on the binomial distribution.²¹ In instances where a study specified a zero occurrence rate of an outcome, the standard error was estimated using a constant continuity correction.²² If a study did not specifically report an outcome, no correction was used, and the study was not included in the analysis of that outcome. Total complications for both analyses were retrieved for only defined postoperative complications (i.e., seroma, hematoma, infection/cellulitis, and flap necrosis). Relative risks were calculated from studies that examined complications in both human acellular dermal matrix and submuscular cohorts specifically. For both analyses, the DerSimonian and Laird random-effects method was used based on interstudy heterogeneity.^{23,24} Heterogeneity was assessed using the Q statistic and the I^2 statistic.²⁵ Small Q statistic p values are indicative of statistically significant heterogeneity, whereas I^2 percentages indicate the amount of heterogeneity be-




Fig. 1. Study attrition diagram. TE/I, tissue expander/implant; HADM, human acellular dermal matrix.

tween studies. For any heterogeneity of statistical significance, sources of heterogeneity were explored by means of an exclusion sensitivity analysis. Funnel plots, Egger's regression test, and Begg's rank correlation test were used to assess publication bias.^{26,27} Statistical analysis was performed using MIX 2.0, Professional Software for Meta-analysis plug-in for Microsoft Excel, version 2.0.1.2 (BiostatXL, 2010), and corroborated with RevMan5.0 (Cochrane Collabo-

ration Information Management System, Oxford, United Kingdom).^{28,29}

RESULTS

Study Characteristics

An English-language PubMed search of MEDLINE identified 879 articles that were eligible for screening. An additional 22 articles were identified through a manual bibliography search,

resulting in a total of 901 citations. Of these, 823 were rejected after a review of titles and abstracts (Fig. 1). After full-text review of the remaining 78 articles, 30 were rejected. All 48 articles that were included were uncontrolled cohort studies. These 48 studies were used to develop pooled summary complication rates. Thirteen studies had information only on human acellular dermal matrix-based reconstructions.^{2,3,5-7,15,18,30-35} Twenty-nine studies had information only on submuscular-based reconstructions.³⁶⁻⁶⁴ Six studies reported complications for both human acellular dermal matrix and submuscular techniques.^{12,13,16,17,65,66} In total, 2037 human acellular dermal matrix reconstructions and 12,847 submuscular reconstructions were included in the meta-analysis (Tables 1 and 2). Both cohorts had similar population demographics (Table 3). Human acellular dermal matrix reconstructions trended toward a higher average intraoperative fill (264.9 ml versus 187.1 ml; $p = 0.10$) when compared with submuscular reconstructions. The six studies used to calculate relative risks represented 877 human acellular dermal matrix reconstructions and 3464 submuscular reconstructions (Table 4).

T1-2
T3

T4

Pooled Complication Rates

There was an increased rate of total complications, 15.4 percent (95 percent confidence interval, 9.3 to 21.4 percent) versus 14.0 percent (95 percent confidence interval, 11.7 to 16.3 percent); seroma, 4.8 percent (95 percent confidence interval, 2.8 to 6.9 percent) versus 3.5 percent (95 percent confidence interval, 2.6 to 4.4 percent); infection, 5.3 percent (95 percent confidence interval, 3.1 to 7.4 percent) versus 4.7 percent (95 percent confidence interval, 3.8 to 5.7 percent); and flap necrosis, 6.9 percent (95 percent confidence interval, 3.6 to 10.2 percent) versus 4.9 percent (95 percent confidence interval, 3.7 to 6.2 percent) in human acellular dermal matrix when compared with submuscular reconstructions (Fig. 2 and Table 5). However, the rate of hematoma was greater in the submuscular cohort (1.5 percent; 95 percent confidence interval, 1.0 to 2.0 percent) than in the human acellular dermal matrix cohort (1.0 percent; 95 percent confidence interval, 0.5 to 1.5 percent), and the rate of reconstructive failure was very similar in both cohorts, 3.8 percent (95 percent confidence interval, 2.3 to 5.4 percent) versus 3.8 percent (95 percent confidence interval, 2.9 to 4.7 percent).

F2,T5

Meta-Analysis of Comparative Studies

There was an increase in the risk of total complications (relative risk, 2.05; 95 percent confidence interval, 1.55 to 2.70), seroma (relative risk, 2.73; 95 percent confidence interval, 1.67 to 4.46), infection (relative risk, 2.47; 95 percent confidence interval, 1.71 to 3.57), and reconstructive failure (relative risk, 2.80; 95 percent confidence interval, 1.76 to 4.45) in the human acellular dermal matrix cohort (Figs. 3 through 5 and Table 6). There was a trend toward increased risk of hematoma (relative risk, 2.06; 95 percent confidence interval, 0.86 to 4.95) and flap necrosis (relative risk, 1.56; 95 percent confidence interval, 0.85 to 2.85) in the human acellular dermal matrix cohort, but the results were not statistically significant.

F3-5,T6

Study Heterogeneity

The majority of pooled complication analyses showed significant heterogeneity (Table 5). Heterogeneity values of the 19 studies in the human acellular dermal matrix cohort (Q statistic, $p < 0.01$; $I^2 = 95.4$ percent) were similar to those of the submuscular cohort (Q statistic, $p < 0.01$; $I^2 = 94.0$ percent). Relative risk analyses were generally more homogeneous, with only total complications and flap necrosis reporting statistically significant heterogeneity (Q statistic, $p = 0.08$; $I^2 = 49.0$ percent; and $p = 0.02$; $I^2 = 68.3$ percent). In analyses of seroma (Q statistic, $p = 0.22$; $I^2 = 29.3$ percent), infection (Q statistic, $p = 0.72$; $I^2 = 0.0$ percent), hematoma (Q statistic, $p = 0.98$; $I^2 = 0.0$ percent), and reconstructive failure (Q statistic, $p = 0.53$; $I^2 = 0.0$ percent), heterogeneity was not significant (Table 6).

Publication Bias

The funnel plot generated to test for publication bias in the relative risk analysis of total complications showed relative symmetry (Fig. 6). Egger's weighted regression and Begg's rank correlation test showed minimal evidence of bias (bias = 0.30; 95 percent confidence interval, -5.96 to 6.57; $p = 0.90$, and Kendall's tau-b = -0.06; $p = 0.85$).

F6

DISCUSSION

The advent of any new technology—and specifically, its integration with a new technique—generates uncertainty about outcomes and questions about benefits vis-à-vis risks. With human acellular dermal matrix-assisted tissue expander breast reconstruction, the potential for improvement in cosmetic and reconstructive outcomes has been promulgated with little consensus on cumula-

Table 1. Characteristics of Study Populations Reporting Human Acellular Dermal Matrix Reconstruction Outcomes

Reference	Patient Characteristics							Complications					
	No. of Reconstructions	Mean Age (yr)	Mean BMI	Previous XRT (%)	Postoperative XRT (%)	Mean Intraoperative Fill (ml)	Mean Follow-Up (mo)	Total Complications	Seroma	Hematoma	Infection	Flap Necrosis	Reconstructive Failure
Antony et al., 2010 ¹⁶	153	44.5	23.8	10.5	9.2	NA	NA	32	11	3	11	7	9
Becker et al., 2009 ³⁰	50	50.9	26.3	0.0	NA	40.0	14.3	2	1	NA	1	NA	NA
Bindingavele et al., 2007 ³¹	65	50.0	NA	NA	12.2	NA	10.0	6	3	1	2	NA	1
Breuing and Colwell, 2007 ¹⁵	67	46.0	NA	7.4	7.4	NA	16.1	2	NA	NA	2	NA	1
Buck et al., 2010 ³	58	53.3	NA	4.0	10.0	213.5	4.6	5	1	1	2	1	2
Chun et al., 2010 ¹²	269	47.0	25.5	8.7	6.5	322.7	NA	131	38	6	24	63	16
Lanier et al., 2010 ⁶⁵	52	51.0	29.8	5.8	5.8	256.0	6.7	31	8	0	15	8	10
Liu et al., 2011 ¹⁷	266	NA	24.9	NA	9.8	187.8	NA	75	19	1	18	37	13
Losken, 2009 ⁷	31	48.0	26.7	NA	25.8	NA	10.2	1	0	NA	0	1	0
Margulies et al., 2005 ³²	50	46.0	NA	NA	NA	NA	7.9	2	NA	NA	1	1	NA
Nahabedian, 2009 ⁶	100	48.2	NA	11.8	18.4	NA	17.0	13	5	NA	5	3	2
Namnoum, 2009 ³³	29	NA	NA	NA	3.4	170.0	21.0	3	1	NA	1	1	NA
Preminger et al., 2008 ⁶⁶	45	NA	NA	NA	NA	223.8	NA	7	3	1	3	NA	NA
Rawlani et al., 2011 ²	121	50.2	NA	2.5	19.0	256.6	10.2	19	2	0	9	8	11
Salzberg et al., 2011 ¹⁸	466	NA	NA	2.4	2.1	412.8	28.9	11	NA	5	1	5	6
Sbitany et al., 2009 ¹³	92	48.6	26.4	NA	12.0	412.5	NA	11	3	NA	8	NA	4
Spear et al., 2008 ⁵	58	50.3	NA	5.2	13.8	242.0	18.1	7	1	NA	4	2	1
Topol et al., 2008 ³⁴	35	NA	NA	NA	NA	NA	9.5	3	NA	NA	3	NA	2
Zienowicz and Karacaoglu, 2007 ³⁵	30	NA	NA	NA	NA	441.0	18.0	6	NA	NA	NA	6	NA

NA, not applicable; XRT, radiotherapy.

Table 2. Characteristics of Study Populations Reporting Submuscular Reconstruction Outcomes

Table 2. Characteristics of Study Populations Reporting Submuscular Reconstruction Outcomes													
Patient Characteristics													
Reference	No. of Reconstructions	Age (yr)	BMI	Mean XRT (%)	Previous XRT (%)	Postoperative XRT (%)	Intraoperative Fill (ml)	Mean Follow-Up (mo)	Total Complications	Seroma	Hematoma	Infection	Flap Necrosis
Alderman et al., 2002 ³⁶	79	47.9	NA	NA	NA	NA	NA	NA	33	NA	26	79	189
Antony et al., 2010 ¹⁶	2910	48.1	26.3	11.1	6.5	15.4	NA	15.5	16	11	18	74	15
Ascherman et al., 2006 ³⁷	123	52.5	NA	3.7	9.5	NA	NA	107	10	8	NA	1	3
Berry et al., 2010 ³⁸	733	50.8	NA	6.8	5.8	17.9	31.0	10	2	NA	NA	7	1
Caffo et al., 2000 ³⁹	108	48.9	NA	NA	NA	NA	NA	10	10	NA	NA	39	92
Castello et al., 2000 ⁴⁰	56	44.0	NA	4.3	5.0	NA	NA	38	19	10	NA	8	NA
Chang et al., 2008 ⁴¹	776	NA	NA	NA	26.2	50.4	NA	33	8	4	7	3	13
Clough et al., 2001 ⁴³	360	47.0	NA	NA	5.0	NA	NA	33	22	2	2	7	1
Chun et al., 2010 ¹²	146	46.2	23.8	5.2	8.2	131.2	NA	NA	NA	NA	NA	NA	NA
Cordiero and McCarthy, 2006 ⁴⁴	1522	NA	NA	5.6	15.3	NA	NA	118	5	10	58	45	33
Collis and Sharpe, 2000 ⁴⁵	197	48.0	NA	NA	NA	32.4	NA	25	18	NA	9	4	5
Delgado et al., 2010 ⁴⁶	400	50.0	26.9	NA	17.1	8.0	NA	5	95	17	NA	23	NA
Erkmen and Stark, 2006 ⁴⁷	70	54.7	24.8	9.3	11.6	74.0	NA	18	8	0	5	9	4
Goodwin et al., 2005 ⁴⁸	691	50.0	24.7	10.7	7.8	74.9	NA	15	6	NA	57	102	5
Lanier et al., 2010 ⁴⁹	75	58.7	34.0	3.7	11.1	328.0	16.0	197	15	38*	NA	57	102
Losken et al., 2010 ⁵⁰	1170	46.0	NA	9.6	NA	NA	NA	6	58	17	3	23	14
McCarthy et al., 2008 ⁵⁰	83	NA	NA	NA	NA	NA	NA	11	58	NA	1	5	5
Nano et al., 2005 ⁵¹	481	NA	NA	NA	NA	NA	NA	3	2	NA	0	10	NA
Padubidhi et al., 2001 ⁵²	65	NA	NA	NA	201.1	2.1	NA	15	5*	6	NA	9	16
Pinsole et al., 2006 ⁵³	45	NA	NA	NA	8.6	271.0	NA	33	6	2	2	13	17
Preminger et al., 2008 ⁵⁴	370	48.0	NA	NA	NA	340.0	NA	39	3	3	6	1	4
Prusic and Cordetro, 2003 ⁵⁵	214	47.0	26.1	13.1	6.5	NA	NA	7	2	0	0	5	0
Radvanovic et al., 2010 ⁵⁵	31	52.0	NA	9.7	26.2	NA	NA	7	3	NA	2	6	17
Saint-Cyr et al., 2010 ⁵⁶	256	47.5	NA	NA	NA	NA	NA	7	3	NA	2	5	0
Sandelin et al., 2004 ⁵⁸	188	51.7	28.2	NA	NA	NA	NA	9	12	3	14	2	NA
Schitany et al., 2009 ⁵⁹	84	47.5	NA	0.0	NA	NA	NA	7	9*	NA	NA	7	NA
Stroock, 2009 ⁶⁰	246	NA	NA	10.4	61.0	NA	NA	17	10	NA	NA	4	4
Talier et al., 2003 ⁶¹	77	46.9	NA	NA	4.8	NA	65.0	10	62	21	4	24	13†
Vandeweyer et al., 2000 ⁶²	124	44.0	23.6	8.3	4.2	138.0	73.4	62	62	21	4	24	13†
Woerdenman et al., 2007 ⁶³	174	44.0	23.6	8.3	4.2	138.0	28.6	168	20	31	61	56†	NA
Woerdenman et al., 2006 ⁶⁴	400	43.0	24.0	6.1	NA	183.0	28.6	168	20	31	61	56†	NA

*Seroma and hematoma combined.													
NA, not applicable; XRT, radiotherapy.													
†Reported as skin problems.													

NA, not applicable; XRT, radiotherapy.
*Seroma and hematoma combined.
†Reported as skin problems.

Table 3. Average Demographic Data

	HADM	Submuscular	<i>p</i>
Mean age, yr	48.8	48.8	0.97
Mean BMI	26.2	26.1	0.94
Smokers, %	11.3	17.6	0.10
Diabetes, %	4.1	2.6	0.21
Previous radiotherapy, %	5.8	6.5	0.64
Postoperative radiotherapy, %	11.1	13.3	0.50
Chemotherapy, %	39.8	43.0	0.67
Mean intraoperative fill, ml	264.9	187.1	0.10
Mean follow-up, mo	13.8	29.9	0.02

HADM, human acellular dermal matrix.

tive risks. The variance in total complication profiles reported in the literature (range, 3.2 to 48.7 percent) suggests the need for pooled analysis.^{2,3,5,7,12,15-18} The aim of our meta-analysis was to focus on complication profiles and perform both a risk analysis and pooled summary estimates using the most recent literature. We did not specifically address the underlying rationale for using human acellular dermal matrix or validate the potential advantages of this technique, as these topics have been discussed elsewhere in the literature.^{2,4,5,6,15}

An important question associated with the use of human acellular dermal matrix is whether or not there is an enhanced infection rate. Our meta-analysis shows that the risk of infection is more than doubled when using human acellular dermal matrix (relative risk, 2.47; 95 percent confidence interval, 1.71 to 3.57). This risk is corroborated by the difference in pooled rates between human acellular dermal matrix and submuscular reconstruction, 5.3 percent (95 percent confidence interval, 3.1 to 7.4 percent) versus 4.7 percent (95 percent confidence interval, 3.8 to 5.7 percent), respectively. This may not be surprising considering that the initial state of the human acellular dermal matrix is that of a pre-vascularization biological material and, as such, another potential foreign body added to the stressed hypovascular milieu of mastectomy flaps. Another concern with human acellular dermal matrix has been the potential for enhanced seroma formation—the rationale being that the biological material may incite an inflammatory response that manifests as seroma (or, alternatively, as the cellulitic mimic termed “red breast syndrome”).^{2,67,68} Indeed, our meta-analysis suggests this to be the case in both the risk analysis and pooled rates (relative risk, 2.73; 95 percent confidence interval, 1.67 to 4.46; and 4.8 percent; 95 percent confidence interval, 2.8 to 6.9 percent, versus 3.5 percent; 95 percent confidence interval, 2.6 to 4.4 percent).

Another complication associated with tissue expander reconstruction is hematoma. The rela-

tive similarity in tissue plane dissection and technique between the human acellular dermal matrix and submuscular cohort would presuppose a hypothesis that hematoma rates would be similar. Indeed, both the meta-analysis and the pooled complication analysis indicate that hematoma rates themselves are not affected by the incorporation of human acellular dermal matrix into the standard submuscular technique.

The rates of mastectomy flap necrosis and human acellular dermal matrix use may be related by intraoperative expansion. With the anatomical constraints of an intact pectoralis major muscle eliminated, the human acellular dermal matrix-based tissue expander reconstruction may allow for greater intraoperative fill as suggested by our analysis (264.9 ml versus 187.1 ml; $p = 0.10$). However, this must be counterbalanced by the possibility of added vascular insult to already compromised mastectomy flaps and the attendant risk of necrosis. Thus, our pooled complication rates suggested a higher rate of flap necrosis in the human acellular dermal matrix cohort, 6.9 percent (95 percent confidence interval, 3.6 to 10.2 percent) versus 4.9 percent (95 percent confidence interval, 3.7 to 6.2 percent). However, the risk analysis did not connote a significant relative risk with this variable (relative risk, 1.56; 95 percent confidence interval, 0.85 to 2.85). Moreover, intraoperative fill showed no apparent correlation with flap necrosis in studies that reported both of these values ($p = 0.56$). Part of the reason for this lack of correlation may be the degree of clinical judgment that factors into expansion—wholesale expansion of prostheses is limited by surgical assessment of mastectomy flaps, which will vary from patient to patient. Finally, with the meta-analysis reflecting an increased total complication profile and relative risk, it is perhaps not surprising that the risk of reconstructive failure or removal of the expander is higher in the human acellular dermal matrix cohort (relative risk, 2.80; 95 percent confidence interval, 1.76 to 4.45). Reducing the incidence of these complications has been discussed by various authors in the literature.^{2,12,16,18} Specifically, strategies that have been used by the senior author (J.Y.S.K.) include judicious intraoperative expansion, careful patient selection based on an intraoperative assessment of flap vascularity, dead space management with the use of quilting sutures, and aggressive irrigation of the breast pocket.

Although beyond the scope of this study, the issue of cost-effectiveness is another important element to consider when performing human acellular dermal matrix-assisted breast reconstruc-

Table 4. Characteristics of Study Populations Comparing Human Acellular Dermal Matrix and Submuscular Outcomes

Reference	No. of Reconstructions	Mean Age (yr)	Mean Intraoperative Fill (ml)	Total Complications	Seroma	Hematoma	Infection	Flap Necrosis	Reconstructive Failure
Antony et al., 2010 ¹⁶	153	44.5	NA	32	11	3	11	7	9
HADM	2910	48.1	NA	341	47	26	79	189	55
Submuscular	269	47.0	322.7	131	38	6	24	63	16
Chun et al., 2010 ¹²	146	46.2	131.2	22	4	2	3	13	1
HADM	52	51.0	256.0	31	8	0	15	8	10
Submuscular	75	50.0	74.0	18	5	0	9	4	4
Liu et al., 2011 ¹⁷	266	48.0	187.8	75	19	1	18	37	13
HADM	204	50.0	74.9	35	8	0	5	22	5
Submuscular	45	NA	223.8	7	3	1	3	NA	NA
Preminger et al., 2008 ⁶⁶	45	NA	201.1	3	2	0	1	NA	NA
Submuscular	92	48.6	412.5	11	3	NA	8	NA	4
Sbitany et al., 2009 ¹³	84	51.7	130.0	9	3	NA	6	NA	3
Submuscular	HADM, human acellular dermal matrix; NA, not applicable.								

Pooled Rate of Complications in HADM Reconstruction




Fig. 2. Cumulative pooled rate of total complications in human acellular dermal matrix (HADM) reconstruction. *Diamonds* indicate the cumulative pooled rate of partial flap loss in individual studies. *Horizontal lines* represent the corresponding 95 percent confidence interval (CI). The *vertical dashed line* represents the cumulative rate estimate.

Table 5. Pooled Complication Rates for Human Acellular Dermal Matrix and Submuscular Patients

HADM	No. of Studies	No. of Reconstructions	Pooled Complication Rate (%)	95% CI (%)	Q Statistic, p	I ² (%)
Total complications	19	2037	15.4	9.3-21.4	<0.01	95.4
Seroma	14	1389	4.8	2.8-6.9	<0.01	71.0
Hematoma	9	1495	1.0	0.5-1.5	0.68	0.0
Infection	18	2007	5.3	3.1-7.4	<0.01	83.9
Flap necrosis	13	1683	6.9	3.6-10.2	<0.01	90.0
Reconstructive failure	14	1833	3.8	2.3-5.4	<0.01	64.0
Submuscular						
Total complications	35	12,847	14.0	11.7-16.3	<0.01	94.0
Seroma	24	9886	3.5	2.6-4.4	<0.01	85.7
Hematoma	22	9547	1.5	1.0-2.0	<0.01	65.7
Infection	35	12,847	4.7	3.8-5.7	<0.01	85.8
Flap necrosis	26	10,818	4.9	3.7-6.2	<0.01	91.2
Reconstructive failure	31	12,144	3.8	2.9-4.7	<0.01	88.3

CI, confidence interval; HADM, human acellular dermal matrix.

tion. Our study does not specifically address this topic; however, each surgeon considering the potential benefits of human acellular dermal matrix must also consider the potential economic burden and aforementioned complication profile as well.

The statistical tools used to analyze the studies were stratified to formal meta-analysis for studies with both human acellular dermal matrix and submuscular cohorts and to cumulative pooled complication summaries for studies with solely human

Total Complications in HADM and Submuscular Reconstruction




Fig. 3. Total complications in human acellular dermal matrix (HADM) and submuscular reconstruction. Size of the solid squares is inversely proportional to the variance of the study estimate. The diamond represents the random-effects relative risk and 95 percent confidence interval (CI). The dashed line represents the overall risk estimate.

Seroma in HADM and Submuscular Reconstruction




Fig. 4. Seroma in human acellular dermal matrix (HADM) and submuscular reconstruction. Size of the solid squares is inversely proportional to the variance of the study estimate. The diamond represents the random-effects relative risk and 95 percent confidence interval (CI). The dashed line represents the overall risk estimate.

acellular dermal matrix or submuscular cohorts. An earlier study performed a noncomparative complication summary (no submuscular cohort) with a similar range of human acellular dermal matrix complications as seen in our analysis, but our specific meta-analysis allows for an introduction of relative risk and a more up-to-date complication profile.⁶⁹ Even with the inclusion of a

more robust analysis, the data are hampered by missingness because of the inconsistent reporting of variables in the literature. This may explain the lack of perfect concordance between the pooled complication rates and relative risk.

Our heterogeneity values for the pooled complications in both cohorts were significant. We explored causes of heterogeneity, which we hy-




Fig. 5. Infection in human acellular dermal matrix (HADM) and submuscular reconstruction. Size of the solid squares is inversely proportional to the variance of the study estimate. The diamond represents the random-effects relative risk and 95 percent confidence interval (CI). The dashed line represents the overall risk estimate.

Table 6. Relative Risks for Human Acellular Dermal Matrix versus Submuscular Reconstruction

	No. of Studies	Relative Risk	95% CI	Q Statistic, p	P (%)
Total complications	6	2.05	1.55–2.70	0.08	49.0
Seroma	6	2.73	1.67–4.46	0.22	29.3
Hematoma	4	2.06	0.86–4.94	0.98	0.0
Infection	6	2.47	1.71–3.57	0.72	0.0
Flap necrosis	4	1.56	0.85–2.85	0.02	68.3
Reconstructive failure	5	2.80	1.76–4.45	0.53	0.0

CI, confidence interval.

pothesize stem from clinical and methodologic diversity between the studies. Rather than ignoring the heterogeneity present in our study by using a fixed-effects model, we incorporated the heterogeneity through the use of the random-effects model. We had similar results calculating relative risk when applying both models, and varying results for pooled complications. As a result, we chose the more conservative, random-effects estimate for both. In addition, when an exclusion sensitivity assessment was performed by removing study population outliers, there was little change in the statistical significance of the majority of our results. Nevertheless, the relative imbalance of cohort sample sizes in human acellular dermal matrix versus submuscular reconstruction ($n = 2037$ versus $n = 12,847$) introduces the possibility of heightened β error when combined with the missingness factor. Another limitation to this study was that the meta-analysis was based primarily on non-

randomized studies. Ideally, a more robust analysis would require level I evidence with randomized controlled trials. Nonrandomized studies tend to be more susceptible to selection bias because of the potential for greater systematic differences between cohorts as compared with randomized controlled trials, which use randomization and blinding to address this bias. The significant heterogeneity seen in our study (as with most meta-analyses including nonrandomized studies) was expected and is likely related to the increased interstudy methodologic diversity characteristic of nonrandomized study inclusion. The impact of this bias can be mitigated by using strong study inclusion criteria and the same random-effects model we used to draw statistical conclusions.

These limitations and assumptions notwithstanding, the pooled complication rates and risk analysis both demonstrate convergent patterns of increased complications in the human acellular




Fig. 6. Funnel plot of log relative risks according to their precision for total complications. The vertical solid line is drawn at the pooled log of relative risk for total complications, and the dashed lines represent the expected 95 percent confidence interval for a given standard error.

dermal matrix cohort. Given that the studies in the human acellular dermal matrix cohort represent the earliest data on a new technique modification, this may not be surprising because some of the reported data will reflect a learning curve phenomenon (whereas the submuscular cohort studies will reflect patient data from a more mature technique that is several decades old). Intrastitutional data will thus vary from early adoption to later proficiency.⁷⁰ Interinstitutional data may differ according to diverse surgical practices, including the use of tumescent solution or the predilection for thinner mastectomy flaps. In total, there were five different acellular matrices used in the meta-analysis, with the majority of cases involving AlloDerm. However, with only one smaller scale matched study specifically analyzing outcomes related to use of differing human acellular dermal matrix technology, no conclusions can be drawn related to differences in human acellular dermal matrix—this would be a key direction for future studies. With respect to differentiating human acellular dermal matrix from submuscular reconstruction, these issues may or may not impact outcomes.

CONCLUSIONS

Human acellular dermal matrix-assisted breast reconstruction is a technique that continues to evolve. Its principal value proposition may be the potential enhancement in cosmesis and amelioration of late or irradiation-induced contracture.^{2,5-7,15,18,71} In this analysis, the average

follow-up time for the human acellular dermal matrix cohort was 13.8 months versus 28.3 months for the submuscular cohort. As more studies with longer follow-up are generated, a more robust meta-analysis will be possible with which to assess these long-term results. Specifically, issues related to revision surgery, capsular contracture, and patient satisfaction could be further elucidated.

Focusing specifically on comparing complication profiles of human acellular dermal matrix and traditional submuscular reconstruction by means of a meta-analysis and pooled summaries, our study demonstrates that human acellular dermal matrix seems to connote a higher complication profile than submuscular reconstruction. Although this may well be an artifact of communal inexperience with a new technique, it is nevertheless an important clinical finding to consider in communicating best practice information to patients. Moreover, future studies that directly link benefits with realized risk will help clarify the utility of this technique.

John Y. S. Kim, M.D.
Division of Plastic and Reconstructive Surgery
Northwestern University
Feinberg School of Medicine
675 North St. Clair Street
Galter Suite 19-250
Chicago, Ill. 60611
jokim@nmh.org

REFERENCES

1. American Society of Plastic Surgeons. Reconstructive breast surgery statistics 2009. Available at: <http://www.plasticsurgery.org>.

Volume 129, Number 1 • Acellular-Assisted Breast Reconstruction

- org/media/statistics/2009Statistics.cfm. Accessed February 16, 2010.
2. Rawlani V, Buck DW II, Johnson SA, Heyer KS, Kim JY. Tissue expander breast reconstruction using prehydrated human acellular dermis. *Ann Plast Surg* 2011;66:593-597.
3. Buck DW II, Heyer K, DiBardino D, Bethke K, Kim JY. Acellular dermis-assisted breast reconstruction with the use of crescentic tissue expansion: A functional cosmetic analysis of 40 consecutive patients. *Aesthet Surg J* 2010;30:194-200.
4. Salberg CA. Nonexpansive immediate breast reconstruction using human acellular tissue matrix graft (AlloDerm). *Ann Plast Surg* 2006;57:1-5.
5. Spear SL, Parikh PM, Reisin E, Menon NG. Acellular dermis-assisted breast reconstruction. *Aesthetic Plast Surg* 2008;32:418-425.
6. Nahabedian MY. AlloDerm performance in the setting of prosthetic breast surgery, infection, and irradiation. *Plast Reconstr Surg* 2009;124:1743-1753.
7. Losken A. Early results using sterilized acellular human dermis (NeoForm) in post-mastectomy tissue expander breast reconstruction. *Plast Reconstr Surg* 2009;123:1654-1658.
8. Kim JYS, Johnson SA, Fine N. Dual-plane tissue expander reconstruction with acellular dermal matrix. In: Losken A, ed. *Innovations in Plastic Surgery*. St Louis: Quality Medical; 2010:33-44.
9. Kim JY, Buck DW II, Kloeters O, Eo S, Jones NF. Reconstruction of a recurrent first dorsal web space defect using acellular dermis. *Hand (N Y.)* 2007;2:240-244.
10. Kim JY, Bullocks JM, Basu CB, et al. Dermal composite flaps reconstructed from acellular dermis: A novel method of neothelial reconstruction. *Plast Reconstr Surg* 2005;115:96-100e.
11. Wainwright DJ. Use of an acellular allograft dermal matrix (AlloDerm) in the management of full-thickness burns. *Burns* 1995;21:243-248.
12. Chun YS, Verma K, Rosen H, et al. Implant-based breast reconstruction using acellular dermal matrix and the risk of post-operative complications. *Plast Reconstr Surg* 2010;125:429-436.
13. Shihani H, Sandeen SN, Amalfi AN, Davenport MS, Langstein HN. Acellular dermis-assisted prosthetic breast reconstruction versus complete submuscular coverage: A head-to-head comparison of outcomes. *Plast Reconstr Surg* 2009;124:1735-1740.
14. Ko JH, Salway DM, Paul BC, Wang EC, Dumanian GA. Soft polypropylene mesh, but not cadaveric dermis, significantly improves outcomes in midline hernia repairs using the components separation technique. *Plast Reconstr Surg* 2009;124:836-847.
15. Breuing KH, Colwell AS. Inferolateral AlloDerm hammock for implant coverage in breast reconstruction. *Ann Plast Surg* 2007;59:250-255.
16. Antony AK, McCarthy CM, Cordeiro PG, et al. Acellular human dermis implantation in 153 immediate two-stage tissue expander breast reconstructions: Determining the incidence and significant predictors of complications. *Plast Reconstr Surg* 2010;125:1606-1614.
17. Lin AS, Kao HK, Reish RG, Herguener CA, May JW Jr, Guo L. Post-operative complications in prosthesis-based breast reconstruction using acellular dermal matrix. *Plast Reconstr Surg* 2011;127:1755-1762.
18. Salberg CA, Ashikari AV, Koch RM, Chabner-Thompson E. An 8-year experience of direct-to-implant immediate breast reconstruction using human acellular dermal matrix (AlloDerm). *Plast Reconstr Surg* 2011;127:514-524.
19. Higgins JT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.0.2 [updated September 2009]. The Cochrane Collaboration, 2009. Available at: <http://www.cochrane-handbook.org>. Accessed January 28, 2011.
20. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: A proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008-2012.
21. Armitage P, Berry G, eds. *Statistical Methods in Medical Research*. 3rd ed. Boston: Blackwell Scientific; 1994.
22. Sweeting MJ, Sutton AJ, Lambert PC. What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data. *Stat Med* 2004;23:1351-1375.
23. Deeks JJ, Altman DG, Bradburn MJ. Statistical methods for examining heterogeneity and combining results from several studies in meta-analysis. In: Egger M, Davey Smith G, Altman DG, eds. *Systematic Reviews in Health Care: Meta-analysis in Context*. London: BMJ; 2001.
24. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-188.
25. Huedo-Medina TB, Sanchez-Meca J, Marin-Martinez F, Botella J. Assessing heterogeneity in meta-analysis: Q statistic or I² index? *Psychol Methods* 2006;11:193-206.
26. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-634.
27. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088-1101.
28. Bax L, Yu LM, Ikeda N, Tsuruta H, Moons KG. Development and validation of MIX: Comprehensive free software for meta-analysis of causal research data. *BMC Med Res Methodol* 2006;6:50.
29. Bax L, Yu LM, Ikeda N, Moons KG. A systematic comparison of software dedicated to meta-analysis of causal studies. *BMC Med Res Methodol* 2007;7:40.
30. Becker S, Saint-Cyr M, Wong C, et al. AlloDerm versus Dermatrix in immediate expander-based breast reconstruction: A preliminary comparison of complication profiles and material compliance. *Plast Reconstr Surg* 2009;123:1-6; discussion 107-108.
31. Bindingavele V, Gaon M, Ora KS, Kulber DA, Lee DJ. Use of acellular cadaveric dermis and tissue expansion in post-mastectomy breast reconstruction. *J Plast Reconstr Aesthet Surg* 2007;60:1214-1218.
32. Margulies AG, Hochberg J, Kepple J, Henry-Tillman RS, Westbrook K, Klimberg VS. Total skin-sparing mastectomy without preservation of the nipple-areola complex. *Am J Surg* 2005;190:907-912.
33. Nannoun JD. Expander/implant reconstruction with AlloDerm: Recent experience. *Plast Reconstr Surg* 2009;124:387-394.
34. Topol BM, Dalton EF, Ponn T, Campbell CJ. Immediate single-stage breast reconstruction using implants and human acellular dermal tissue matrix with adjustment of the lower pole of the breast to reduce unwanted lift. *Ann Plast Surg* 2008;61:494-499.
35. Zienowicz RJ, Karacaoglu E. Implant-based breast reconstruction with allograft. *Plast Reconstr Surg* 2007;120:373-381.
36. Alderman AK, Wilkins EG, Kim HM, Lowery JC. Complications in postmastectomy breast reconstruction: Two-year results of the Michigan Breast Reconstruction Outcome Study. *Plast Reconstr Surg* 2002;109:2265-2274.
37. Ascherman JA, Hanasano MM, Newman MI, Hughes DB. Implant reconstruction in breast cancer patients treated with radiation therapy. *Plast Reconstr Surg* 2006;117:359-365.
38. Berry T, Brooks S, Sydow N, et al. Complication rates of radiation on tissue expander and autologous tissue breast reconstruction. *Ann Surg Oncol* 2010;17:202-210.

39. Caffo O, Cazzolli D, Scalet A, et al. Concurrent adjuvant chemotherapy and immediate breast reconstruction with skin expanders after mastectomy for breast cancer. *Breast Cancer Res Treat.* 2000;60:267-275.
40. Castelló JR, Garro L, Nájera A, Mirelis E, Sánchez-Olaso A, Barros J. Immediate breast reconstruction in two stages using anatomic tissue expansion. *Scand J Plast Reconstr Surg Hand Surg.* 2000;34:167-171.
41. Chang DW, Barnea Y, Robb GL. Effects of an autologous flap combined with an implant for breast reconstruction: An evaluation of 1000 consecutive reconstructions of previously irradiated breasts. *Plast Reconstr Surg.* 2008;122:356-362.
42. Cicchetti S, Leone MS, Franchelli S, Santi PL. One-stage breast reconstruction using McGhan style 150 bi-dimensional expanders: A review of 107 implants with six years experience. *J Plast Reconstr Aesthet Surg.* 2006;59:1037-1042.
43. Clough KB, O'Donoghue JM, Fitoussi AD, Nos C, Falcou MC. Prospective evaluation of late cosmetic results following breast reconstruction: I. Implant reconstruction. *Plast Reconstr Surg.* 2001;107:1702-1709.
44. Cordeiro PG, McCarthy CM. A single surgeon's 12-year experience with tissue expander/implant breast reconstruction: Part I. A prospective analysis of early complications. *Plast Reconstr Surg.* 2006;118:825-831.
45. Collis N, Sharpe DT. Breast reconstruction by tissue expansion: A retrospective technical review of 197 two-stage delayed reconstructions following mastectomy for malignant breast disease in 189 patients. *Br J Plast Surg.* 2000;53:37-41.
46. Delgado JF, García-Guillarte RF, Palazuelo MR, Mendez JI, Pérez CC. Immediate breast reconstruction with direct, anatomic, gel-cohesive, extra-projection prosthesis: 400 cases. *Plast Reconstr Surg.* 2010;125:1599-1605.
47. Eriksen C, Stark B. Early experience with the crescent expander in immediate and delayed breast reconstruction. *Scand J Plast Reconstr Surg Hand Surg.* 2006;40:82-88.
48. Goodwin SJ, McCarthy CM, Pusic AL, et al. Complications in smokers after postmastectomy tissue expander/implant breast reconstruction. *Ann Plast Surg.* 2005;55:16-19; discussion 19-20.
49. Losken A, Collins BA, Carlson GW. Dual-plane prosthetic reconstruction using the modified wise pattern mastectomy and fasciocutaneous flap in women with macromastia. *Plast Reconstr Surg.* 2010;126:731-738.
50. McCarthy CM, Mehrara BJ, Riedel E, et al. Predicting complications following expander/implant breast reconstruction: An outcomes analysis based on preoperative clinical risk. *Plast Reconstr Surg.* 2008;121:1886-1892.
51. Nano MT, Gill PG, Kollias J, Bochner MA, Carter N, Winefield HR. Qualitative assessment of breast reconstruction in a specialist breast unit. *ANZ J Surg.* 2005;75:445-453; discussion 371-372.
52. Padubidri AN, Yetman R, Browne E, et al. Complications of postmastectomy breast reconstructions in smokers, ex-smokers, and nonsmokers. *Plast Reconstr Surg.* 2001;107:342-349; discussion 350-351.
53. Pinsolle V, Grinfeder C, Mathoulin-Pelissier S, Faucher A. Complications analysis of 266 immediate breast reconstructions. *J Plast Reconstr Aesthet Surg.* 2006;59:1017-1024.
54. Pusic AL, Cordeiro PG. An accelerated approach to tissue expansion for breast reconstruction: Experience with intraoperative and rapid postoperative expansion in 370 reconstructions. *Plast Reconstr Surg.* 2003;111:1871-1875.
55. Radovanovic Z, Radovanovic D, Golubovic A, Iakovic-Kapic T, Bokorov B, Mandic A. Early complications after nipple-sparing mastectomy and immediate breast reconstruction with silicone prosthesis: Results of 214 procedures. *Scand J Surg.* 2010;99:115-118.
56. Saint-Cyr M, Dauwe P, Wong C, Thakar H, Nagarkar P, Rohrich RJ. Use of the serratus anterior fascia flap for expander coverage in breast reconstruction. *Plast Reconstr Surg.* 2010;125:1057-1064.
57. Salgarello M, Barone-Adesi L, Terribile D, Masetti R. Update on one-stage immediate breast reconstruction with definitive prosthesis after sparing mastectomies. *Breast.* 2011;20:7-14.
58. Sandelin K, Wickman M, Billgren AM. Oncological outcome after immediate breast reconstruction for invasive breast cancer: A long-term study. *Breast.* 2004;13:210-218.
59. Scuderi N, Alfano C, Campus GV, et al. Multicenter study on breast reconstruction outcome using Becker implants. *Aesthetic Plast Surg.* 2011;35:66-72.
60. Strock LL. Two-stage expander implant reconstruction: Recent experience. *Plast Reconstr Surg.* 2009;124:1429-1436.
61. Tallet AV, Salem N, Moutardier V, et al. Radiotherapy and immediate two-stage breast reconstruction with a tissue expander and implant: Complications and esthetic results. *Int J Radiat Oncol Biol Phys.* 2003;57:136-142.
62. Vandeweyer E, Deraemaeker R, Nogaert JM, Hertens D. Immediate breast reconstruction with implants and adjuvant chemotherapy: A good option? *Acta Chir Belg.* 2003;103:98-101.
63. Woerdeman LA, Hage JJ, Hofland MM, Rutgers EJ. A prospective assessment of surgical risk factors in 400 cases of skin-sparing mastectomy and immediate breast reconstruction with implants to establish selection criteria. *Plast Reconstr Surg.* 2007;119:455-463.
64. Woerdeman LA, Hage JJ, Smeulders MJ, Rutgers EJ, van der Horst CM. Skin-sparing mastectomy and immediate breast reconstruction by use of implants: An assessment of risk factors for complications and cancer control in 120 patients. *Plast Reconstr Surg.* 2006;118:321-330; discussion 331-332.
65. Lanier ST, Wang ED, Chen JJ, et al. The effect of acellular dermal matrix use on complication rates in tissue expander/implant breast reconstruction. *Ann Plast Surg.* 2010;64:674-678.
66. Preminger BA, McCarthy CM, Hu QY, Mehrara BJ, Disa JJ. The influence of AlloDerm on expander dynamics and complications in the setting of immediate tissue expander/implant reconstruction: A matched-cohort study. *Ann Plast Surg.* 2008;60:510-513.
67. Heyer K, Buck DW II, Kato C, Khan SA, Alam M, Kim JY. Reversed acellular dermis: Failure of graft incorporation in primary tissue expander breast reconstruction resulting in recurrent breast cellulitis. *Plast Reconstr Surg.* 2010;125:666-686.
68. Kim JYS, Johnson SA, Fine N. Dual-plane tissue expander reconstruction with acellular dermal matrix. In: Losken A, ed. *Innovations in Plastic Surgery*. St. Louis: Quality Medical; 2010:33-44.
69. Newman MJ, Swartz KA, Samson MC, Mahoney CB, Diab K. The true incidence of near-term postoperative complications in prosthetic breast reconstruction utilizing human acellular dermal matrices: A meta-analysis. *Aesthetic Plast Surg.* 2011;35:100-106.
70. Carry MJ, Chan R, Huckman R, Snow D, Orgill DP. A detailed analysis of the reduction mammoplasty learning curve: A statistical process model for approaching surgical performance improvement. *Plast Reconstr Surg.* 2009;124:706-714.
71. Spear SL, Seruya M, Clemens MW, Teitelbaum S, Nahabedian MY. Acellular dermal matrix for the treatment and prevention of implant-associated breast deformities. *Plast Reconstr Surg.* 2011;127:1047-1058.